

# Epidemiology of subtle, typical, cystic, and deep endometriosis: a systematic review

Philippe R. Koninckx<sup>1,2,3,4,5,6</sup> · Anastasia Ussia<sup>3,5</sup> · Jörg Keckstein<sup>7,8</sup> · Arnaud Wattiez<sup>9</sup> · Leila Adamyan<sup>4</sup>

Received: 22 June 2016 / Accepted: 19 July 2016 / Published online: 13 August 2016  
© Springer-Verlag Berlin Heidelberg 2016

**Abstract** Endometriosis is known as a cause of pelvic pain and infertility. The epidemiology of endometriosis is important since the prevalence and severity of endometriosis might be linked to pollution and to our modern lifestyle, comprising food intake, chemical disruptors, postponement of the first pregnancy, and stress, as indicated by “a career women’s disease.” Epidemiological data based upon hospital discharge records should be viewed with caution. Indeed, that subtle lesions are considered pathology causes a major increase in prevalence, while the laparoscopic recognition and histological confirmation of subtle and typical lesions vary with the expertise and interest

of the surgeon. The epidemiological data published by surgical groups on severe forms, as cystic and deep endometriosis, have a referral bias and lack the numbers required for meaningful statistics. Fundamental to understanding epidemiology of endometriosis is that it is unclear that all presentations of endometriosis constitute one disease. We therefore performed a systematic review of the incidences and severity of subtle, typical, cystic, and deep endometriosis lesions separately. The only data found were that severe endometriosis carries a greater hereditary risk and that the prevalence of subtle endometriosis decreases with age whereas the prevalences of typical, cystic, and deep endometriosis increase with age. Surgeons that witnessed over the last 20 years in over 1000 interventions each the evolution of deep endometriosis, however, had a strong impression that severity and prevalence of deep endometriosis are increasing. In conclusion, there are no solid epidemiologic data of each type of endometriosis separately. With all restrictions imposed by the referral bias and by a clinical impression, the consistency of the observation of deep endometriosis surgeons should be a reason for concern. The investigation of the epidemiology of deep endometriosis is suggested since it is relevant because it is clinically severe pathologic and feasible since it can be done from hospital-based records. Indeed, most women will ultimately have surgery with solid information on severity and an unbiased diagnosis if defined as “adenomyosis externa.”

## Precis

In the absence of data, the clinical impression of deep endometriosis surgeons that severity and prevalence of deep endometriosis are increasing is a reason for concern.

✉ Philippe R. Koninckx  
Pkoninckx@gmail.com

- <sup>1</sup> KU Leuven, Leuven, Belgium
- <sup>2</sup> University of Oxford, Oxford, UK
- <sup>3</sup> Università Cattolica, Rome, Italy
- <sup>4</sup> Department of Operative Gynecology, The V.I. Kulakov Research Center for Obstetrics, Gynecology and Perinatology, Ministry of Health of the Russian Federation and Dpt of Reproductive Medicine and Surgery, Moscow State University of Medicine and Dentistry, Moscow, Russia
- <sup>5</sup> Gruppo Italo Belga, Villa del Rosario, Rome, Italy
- <sup>6</sup> Vuilenbos 2, 3360 Bierbeek, Belgium
- <sup>7</sup> Landeskrankenhaus Villach, Villach, Austria
- <sup>8</sup> University Ulm, Ulm, Germany
- <sup>9</sup> Prof OBGYN, University of Strassbourg, France and Latiffa Hospital, Dubai, United Arab Emirates

**Keywords** Endometriosis · Epidemiology · Prevalence · Severity

## Introduction


Understanding the epidemiology of endometriosis is important since the prevalence and severity of endometriosis are eventually linked to pollution such as dioxin [1], PCB [2], and radioactivity

[3]. It has been linked to our diet and our fat consumption and to our style of living [4–6] with postponement of the first pregnancy and stress, both summarized in a “career women’s disease.” Most important is that understanding if and why prevalence and severity are increasing would permit prevention.

Endometriosis is a cause of pain and infertility and considered progressive and recurrent. This generates fear and anxiousness as reflected by the 22,398 peer-reviewed articles on PubMed (February 1, 2016), with an exponential increase since 2000 (Fig. 1), in the number of specific congresses, in the patient support groups, and in discussion groups on social media. It moreover emphasizes the associated costs of suffering, absence from work and of associated medical treatments, infertility treatments, and often repetitive surgeries.

The excellent articles on the prevalence of endometriosis [4–6] and of the analysis of the relationship with potential causal factors and with other diseases [7–12] unfortunately are not conclusive. These epidemiological studies have methodological problems which are the definition of endometriosis which permits to include asymptomatic women, the selection bias induced by the need of a surgical diagnosis, and the specific difficulties to perform cohort or case-control studies [6, 13]. The literature on epidemiology of endometriosis becomes even more confusing when the biases in observations and definitions over time are considered. Until the introduction of laparoscopy, endometriosis was limited to severe lesions as described in the beginning of the century [14–20] and to lesions found accidentally during surgery in the pelvis [21, 22] or in the umbilicus [23]. With the introduction of endoscopy in the late 1960s, the apparent prevalence increased since black-pigmented lesions or typical endometriosis became a frequent observation in women with pain and/or infertility [24]. In the mid-1980s, the prevalence almost doubled by the recognition of non-pigmented or subtle lesions [25–28]. From the 1990s onwards, the recognition of deep endometriosis increased the apparent prevalence even further.

Clinically, epidemiology of endometriosis is even limited since the laparoscopy required to make the diagnosis limits repetitive observations over time, since recognition varies with the training and the interest of the surgeon, and since



**Fig. 1** Number of articles on endometriosis listed on PubMed. Indicated are blocks of 5 years since 1975

confirmation by pathology is only 50 % [29] to 76 % [30] and even less for subtle lesions.

The pathophysiology and the natural history of the disease are unknown, and it remains unclear whether endometriosis is one or several different pathologies, whether all subtle and microscopical lesions are pathology, and whether it is a progressive and recurrent disease. Since clinically severe lesions as cystic ovarian and deep endometriosis are more important, we decided to review the data on epidemiology of subtle, typical, cystic, and deep endometriosis separately.

## Materials and methods

### Epidemiology

PubMed was screened for the keywords epidemiology OR prevalence OR incidence AND endometriosis generating 2656 articles. When instead of endometriosis “subtle endometriosis” OR “non-colored endometriosis” OR “microscopical endometriosis” was used, only 14 articles were found; “typical endometriosis” generated 5 articles, “endometrioma” OR “cystic endometriosis” OR “cystic ovarian endometriosis” 123, and “deep endometriosis” 37 articles. These articles for subtle, typical, cystic, and deep were subsequently searched manually. A distinction between subtle and typical lesions was made in two articles only. All other articles described the incidence of subtle and typical lesions together. Not a single article permitted to evaluate epidemiology over time for subtle/typical lesions, for cystic ovarian endometriosis, or for deep endometriosis. We therefore hand searched the 1409 articles found by “endometriosis” AND “epidemiology/incidence/prevalence” published between 2005 and 2016 without finding a single additional useful article.

### Pathophysiology and definitions of endometriosis

In order to evaluate whether endometriosis should be considered one progressive disease with different degrees of severity or whether the different presentations of endometriosis should better be considered separately, we searched PubMed with the keywords endometriosis AND “pathophysiology” OR “natural history”. The 400 articles over the last 14 years were hand searched. These articles described associated biochemical, immunological, inflammatory, endometrial, genetic, peritoneal fluid, or innervation changes or associations with waste products. Some were opinion papers. None of these articles nor any of the previous articles could explain the pathophysiology and natural history leading to the various clinical manifestations. How to interpret microscopical and on subtle or non-pigmented lesions caused most controversy.

## Results

### One or several diseases?

Endometriosis was defined more than 100 years ago as “endometrial glands and stroma outside the uterus” since microscopy was the only tool available at that moment [31, 32]. The clinical content of that definition changed over time. In the beginning of the twentieth century [14–18, 33], adenomyosis externa was described. The name endometriosis [19, 20] was introduced by Sampson when he described glands and stroma in ovarian “chocolate cysts.” In 1940, he described the theory of retrograde menstruation and implantation [34], and in 1942, Gruenwald described the theory of metaplasia [35]. Only with the introduction of laparoscopy, we realized how frequent typical lesions were.

Following the observations that retrograde menstruation [36, 37] with viable cells [38] occurs in almost all women, it was logic to look for early lesions after implantation. In 1985, macroscopically visible non-pigmented lesions without surrounding sclerosis [25, 33] later called subtle lesions [26] were described. Somewhat later, minimal lesions by scanning electron microscopy [39, 40] and microscopical lesions in normal looking peritoneum were found in some 6 to 13 % of women without and with endometriosis, respectively [41–44]. In the early 1990s, some of these subtle lesions were described to disappear spontaneously called remodeling.

*What is the significance of subtle lesions and of microscopical endometriosis?*

Subtle lesions are considered as the early lesions after implantation, and it was postulated but never proven that these lesions will progress to more severe lesions in some women. However, since subtle lesions were found in up to 90 % of women with pain and or infertility [45], since progression was never observed, and in the absence of evidence that subtle lesions are a cause of infertility or pain, it was suggested and it remains debated that “subtle lesions are a normal condition occurring at least intermittently in most women” [46–48]. An indirect argument to consider subtle and typical lesions differently is that the luteinized unruptured follicle syndrome, a cause of infertility, is associated with typical lesions and not with subtle lesions [49].

Also for peritoneal microscopical endometriosis, there is no evidence for progression or for causing pain or infertility. Recently, microscopical lesions were also found in the perirectal lymph nodes of 18 to 20 % of women with deep endometriosis [50]. Yet, a clinical pathology of this has not yet been described. Similarly, in bowel resections for deep endometriosis, microscopic nest of glands and stroma can be found at least up to 5 cm from the nodule [51, 52]. That recurrence rates following bowel resections and following a more

conservative local resection for deep endometriosis are not manifestly different is another argument to postulate that these microscopical nests of glands and stroma are not always pathology [53, 54].

In the absence of evidence that all microscopical endometriosis and subtle lesions will progress or cause pain or infertility, these lesions should not be considered disease in all women. This does not exclude that some of these lesions might progress, but today, we cannot distinguish between the majority of these lesions that have no clinical importance and may disappear spontaneously and those that will develop into endometriosis lesions causing pain and infertility [53]. Therefore, subtle and microscopical lesions should at least be considered separately until we understand their significance.

### *Pathophysiology and natural history*

None of the pathogenetic theories based upon retrograde menstruation [34], coelomic metaplasia [35], or Müllerian remnants can explain all the different types of endometriosis [55, 56]. With the observation of an increased prevalence of endometriosis in women with obstructed outflow [57–60], the never proven hypothesis is that after implantation or metaplasia, these “normal endometrial cells” will progress inevitably into more severe lesions as typical, cystic ovarian, or deep endometriosis. This theory, however, cannot explain why progression occurs in some women only and why typical, or cystic, or deep lesions develop [55]. In order to fill this gap, the endometriotic disease theory [61] postulated that progression started with a cellular incident and that the type of cellular incident or a mutation would determine progression into typical, cystic, or deep endometriosis, similar to what is seen in most benign tumors. This would explain that endometriosis is hereditary [62–69] and that cystic and deep lesions are clonal in origin [33, 70, 71]. It also explains that dioxin and total body radiation, both acting at the level of the genome, might be causally related to endometriosis [72]. Similarly, the many differences in the endometrium of women with or without endometriosis could thus be interpreted as a sign of susceptibility rather than as the consequence of endometriosis.

Whether these endometriotic cells are normal or not, they develop in the environment of the peritoneal cavity or of the ovary. Peritoneal fluid is a specific microenvironment with concentrations of steroid hormones, cytokines, immunology [73, 74], angiogenetic factors, and many others, that are much different from plasma [75]. In endometriosis patients, many of these are different, but it remains questionable whether this is the cause or the consequence of endometriosis. That the decreased natural killer cell activity in plasma [76] and in peritoneal fluid [77] remains decreased after surgical excision suggests that this might be causally related [78]. Also, the intra-ovarian hormone concentrations are much higher, and

also immunology is different, as evidenced by the preferential implantation of tumor cells as in the Krükenberg tumor. Lastly, especially considering also adenomyosis, pale cells [79] and the role of the junctional zone should be considered. The latter broadens the view to immunology of pregnancy, induction of the physiologic changes of the spiral arteries [80–85], and pre-eclampsia. The tumor-like development of endometrial cells on the chicken allantoic membrane [86] remains intriguing.

*Subtle, typical, cystic, and deep endometriosis should be considered separately*

In the absence of a clear model for pathophysiology and natural history of endometriosis and since not all glands and stroma outside the uterus constitute clinical pathology, and considering in addition adenomyosis and Müllerianosis and extra-pelvic localizations, it seems wise to consider subtle, typical, cystic ovarian, and deep lesions separately, at least until proven that it is one disease.

Typical, cystic, and deep endometriosis should not be considered a recurrent and progressive disease. Recurrence rates of each type of endometriosis are different. Following surgical excision, recurrence rates are estimated around 20 % for typical lesions, around 7 % after stripping and 20 % after superficial destruction of cystic ovarian endometriosis [87, 88], and very low (less than 5 % bowel recurrences) after excision of deep endometriosis. Progression from one type to another has not been observed. On the contrary, almost all three types can be considered clinically end stages, which at the moment of diagnosis are no longer progressive and which by pathology have a non-active burnt out aspect. Cystic ovarian endometriosis does not always increase in volume, and low rectovaginal deep lesions followed clinically over time remain mostly unchanged at least over a few years. Clinically, it seems that after a period of growth, progression will stop with surrounding fibrosis as the remnant of a previous inflammation. This is not in contradiction with the rare cases, in our experience some 10 in 3000 deep endometriosis nodules (PK, AU) that can be fast progressive. This highlights heterogeneity as supported by the fact that during pregnancy, some deep lesions behave differently and can cause spontaneous bowel or bladder ruptures [89].

### **Accuracy of the diagnosis of endometriosis lesions and epidemiology**

The recognition and the confirmation by pathology of subtle lesions vary with the skill, interest, and expertise of the surgeon and the pathologist. Rarely considered is the bias caused by the surgeon, who scrutinizes women with pain or infertility in order to have a diagnosis. Even for typical lesions, the histologic confirmation is only 50 % [29] to 76 % [30].

Endometriosis of the diaphragm is underreported since not all surgeons will systematically inspect the diaphragm in steep anti-Trendelenburg with a 30° scope. Most important is that the recognition in the mid-1980s of non-pigmented lesions or subtle lesions [25, 26, 33] almost doubled the prevalence of endometriosis. Before that period, women with minimal and mild endometriosis were women with typical lesions, whereas thereafter, they became women with subtle lesions and/or typical lesions.

The prevalence of cystic ovarian endometriosis is biased by a variable inclusion of cystic corpora lutea, estimated up to 30 %. To distinguish between them can be difficult by ultrasound and also during surgery. CA125 in chocolate fluid could be used, but a rapid test during surgery is not available [90]. Cystic ovarian endometriosis is strongly associated with adhesions [45], and a chocolate cyst without adhesions has a high probability of being a cystic corpus luteum. The inspection of the inside of the cyst by ovarioscopy [91] was suggested as not conclusive. Moreover, it is often unclear whether diagnosis was confirmed by pathology or “compatible with endometriosis,” without a positive identification of endometrial glands and stroma [92, 93]. The recent diagnosis of small cystic ovarian endometriosis by ultrasound [94, 95] increased the prevalence, as is evidenced by the actual discussion whether small cysts should be left alone or should be treated by laparoscopy or trans hydroculdoscopy [96], or whether it is preferable to proceed to IVF without surgery.

The clinical diagnosis of deep endometriosis is unreliable since in only 50 % of the larger nodules, the diagnosis is made by clinical exam [45, 97]. The diagnostic accuracy of MRI and of ultrasound hardly exceeds 90 % sensitivity and specificity, and it is unclear whether imaging is really helpful in the diagnosis of smaller lesions. Even the surgical diagnosis is biased by a variable indication for surgery, a variable recognition during surgery, a variable referral bias, and the absence of an unanimously accepted definition. Deep endometriosis is partially underreported, since smaller lesions are easily missed especially in the sigmoid. Deep endometriosis is suspected to be partially over-reported since deep endometriosis is considered the ultimate surgical challenge in gynecology; many centers interpret broadly the diagnosis of deep endometriosis in order to belong to the club of hospitals that perform this surgery. Underdiagnosed is deep endometriosis around the sciatic nerve and around the sacral nerve roots because of the surgical risk. It would be preferable to use as definition adenomyosis externa [98, 99], since with the definition of more than 5 mm under the peritoneal surface [99], many typical lesions can be classified as deep endometriosis.

Considering the prevalences of the lesions, women with the diagnosis of endometriosis in hospital discharge records mainly have typical endometriosis and/or subtle endometriosis, some cystic endometriosis, and a few deep endometriosis. Confirmation by pathology will be variable [100, 101].

## Epidemiology of endometriosis lesions

### Epidemiology and age and heredity

With all restrictions imposed by the inherent selection bias in Leuven in 1991, in a group of 900 women with pain and/or infertility, 49 % had subtle lesions, 29 % had typical lesions, 31 % had cystic ovarian endometriosis, and 18 % had deep endometriosis. Although the total incidence remained constant with age at some 71 %, subtle lesions significantly decreased with age whereas typical cystic and deep lesions increased with age [45] (Fig. 2).

Endometriosis is a hereditary disease [62–68, 102] as evidenced by the seven times higher prevalences in first-degree relatives and the high associations in monozygotic. Genome-wide association studies could identify common genetic variants [103–107]. Also, severity and early onset seem hereditary.


### Subtle lesions

Subtle endometriosis will be found in most women when scrutinized during laparoscopy. They are expected to be found in all women if laparoscopy could be done repetitively.

In women with pain and or infertility, the incidence of subtle lesions decreases with age [45, 108]. In adolescents with pain endometriosis (probably mainly subtle lesions), the incidence of subtle lesions was found in 40 % [109] and 70 % [110]. Also in women without symptoms, endometriosis was found in some 40 % [43, 109]. Not a single study permits to evaluate specifically the epidemiology of subtle endometriosis over time or to evaluate an association with early menarche, short cycles, abundant or painful periods, subfertility, canalization defects of the cervix, race, dioxin, total body radiation, or any other factor. There is no evidence that prevalence of subtle lesions is increasing over the last decades.

### Typical lesions

Ignoring the bias of a variable inclusion of subtle lesions, typical endometriosis was reported in up to 5 % in asymptomatic women, e.g., in 4 % in tubal ligation [29], in 0.16 % in white females [111], and in 2.2 % in a large study from Norway [112]. In the latter study, the risk increased with early menarche, frequent menstruations, pelvic pain, infertility, and nulliparity. In women with infertility stage II, endometriosis (mainly typical lesions) was more frequent [113] when



**Fig. 2** Prevalence (red) of subtle, typical, and deep endometriosis in women with infertility ( $n = 1297$ ), pain ( $n = 918$ ), and infertility and pain ( $n = 267$ ). Subtle endometriosis decreases with age, whereas

typical, cystic, and deep endometrioses increase with age. The total prevalence (yellow) remains unchanged (from [45])



the partner was normal (5.7 %) than when azoospermic (3.3 %). In women with pain, higher incidences were found. In teenagers with intractable dysmenorrhea, 50 % had endometriosis [114], and in women with pain and infertility, prevalences of 40 to 70 % were reported, with a mean of 33 % after meta-analysis [115].

No studies permitted to judge the prevalence of typical lesions over the last decades.

In studies investigating the causes of endometriosis, typical and subtle endometrioses were the predominant lesions. Racial differences were repeatedly suggested, but it remains unclear whether black women have lower and Orientals higher rates than Caucasian. That endometriosis increases with earlier menarche and abundant retrograde menstruation is not substantiated. The baboon data describing an increased prevalence of endometriosis following uterine outflow obstruction are not retained since in recent studies, retrograde menstruation was not found [116]. Dioxin pollution has been suggested to be causally related to endometriosis [1, 117] but in the human evidence is scanty [72, 106, 118]. Following the Seveso accident with severe dioxin pollution, the prevalence doubled although not significantly [119]. Although breast-fed girls have been exposed to dioxins in mother milk, they have a lower incidence of endometriosis in adult life [120]. There are no data linking total body radiation to an increased prevalence of endometriosis in the human. It is unclear whether the immune system is causally related to the development of endometriosis notwithstanding the many changes in the immune system [73, 121, 122] and the decreased natural killer cell activity in plasma and in peritoneal fluid [76–78, 123–128]. Indeed, endometriosis prevalence seems not to be obviously affected by chronic immunosuppression, e.g., in transplant patients, nor by smoking affecting NK activity, nor by caffeine or alcohol. It is unclear whether stress is causally related to endometriosis. The only data are the associations between typical endometriosis and a higher trait anxiety [129–131]. The much lower steroid hormone concentrations in peritoneal fluid of women with the LUF syndrome moreover might affect the implantation and/or development of endometriosis [36]. That endometriosis is a career women's disease suggests a causal relationship with stress. It however also points to the delay of first pregnancy with the inevitable increase of infertility with age, an increase in laparoscopies, and thus a higher documented prevalence of endometriosis. This is consistent with a decreasing prevalence with increasing parity [132]. Oral contraception use has been reported to be associated with a slightly decreased prevalence [133].

### Cystic lesions

There is no evidence that the prevalence of cystic ovarian endometriosis is increasing. There is no evidence in

the human for an association between cystic ovarian endometriosis and pollution or lifestyle. The relationship with ovarian cancer is unclear and beyond the scope of this manuscript.

### Deep lesions

The recognition of smaller deep endometriosis lesions only started in the 1990s, and all reports do have a referral bias. An indirect estimation of the prevalence in the Belgian population can be derived from a total of some 500 interventions/year in a population of 10 M (PK and J. Donnez) in the early 1990s, a period that deep endometriosis surgery was systematically referred. Assuming a reproductive life span of 30 years, and a diagnosis and surgery in some half of the women, prevalence in population ranges between 0.2 and 0.5 %. In women with pain and infertility, prevalence was estimated between 3 and 10 % in Belgium as derived from the 10 to 20 % incidence of deep endometriosis in Leuven from 1988 to 1991. Indeed, during that period, referrals were only for pain or infertility since deep endometriosis was not yet well known.

There are no valid data indicating that the prevalence of deep endometriosis is increasing or that deep endometriosis is caused by pollution or lifestyle. A clinical impression of epidemiology of deep endometriosis was asked to internationally recognize deep endometriosis surgeons who performed surgery over the last 20 years and who each performed between 1000 and 3000 interventions of deep endometriosis. With all restrictions imposed by the referral bias each was well aware of, AW, LA, JD, JK, AU, AS, and PK (authors and acknowledged) have the strong impression that severity and probably prevalence of deep endometriosis are increasing; for CK, it is unclear, whereas CN and his brothers consider it a referral bias. Of specific interest is that in the 1990s, surgery for deep endometriosis in the south of Italy was rare, anyway much lower than in Belgium, which contrasts with the high incidence of severe endometriosis we actually observe (AU, AW, PK).

### Adenomyosis, Müllerianosis, peritoneal pockets, and stromatosis

These diseases which are often considered as variants of endometriosis will not be discussed. Stromatosis does not contain endometrial cells, while Müllerianosis is a too rare condition to discuss prevalence. The pathophysiology of adenomyosis is even more enigmatic than endometriosis. Several forms exist, and prevalences vary with the diagnostic method.

## Discussion

Epidemiology of endometriosis is important since endometriosis can be the cause of almost any gynecologic complaint such as infertility, pain, or bleeding problems up to fatigue and sexual problems. Therefore, women and physicians at least consider endometriosis for almost any gynecological complaint. In order to avoid surgery, many women with pain but without endometriosis are treated for endometriosis and live with the idea that they have endometriosis. They all are worried since they fear that endometriosis will cause infertility, that endometriosis is progressive with a high risk of hysterectomy later in life, that there is no effective treatment, and that surgery can be dangerous. After the millennium, awareness grew for pollution, radioactivity, and global warming, which all could be linked to endometriosis. In addition, over the last decades, we witnessed a progressive delay of the first pregnancy, while age, infertility, and endometriosis are correlated. All this, especially in a period of social media, generates a climate of fear of endometriosis. The questions are whether the prevalence of endometriosis and especially severe endometriosis is increasing in our western population and what causes an eventual increase. Asking the question is suggesting the answer as we spontaneously suspect pollution, radioactivity, our lifestyle, and food industry. Since in addition endometriosis has been linked to cancer, it is not surprising that this is fueling fear and concern as expressed by the many support and discussion groups and the explosion of the number of articles [134].

The epidemiology of endometriosis is not clear despite the many excellent articles. Besides the difficulties in collecting epidemiologic data for endometriosis [13, 100, 101] and the difficulties in recognition of endometriosis and in ascertaining the quality of the diagnoses of each form of endometriosis, a key problem is the definition of endometriosis and our prevailing concept of pathophysiology and natural history. With the definition “glands and stroma outside the uterus,” the widespread belief is that all forms of endometriosis fitting that definition are one disease, progression after implantation is considered inevitable, and endometriosis is considered a progressive and recurrent disease.

As discussed, this concept can be challenged. Since glands and stroma outside the uterus do not mean pathology by definition, the word endometriosis is confusing since it is used for both normal and pathologic conditions. It indeed was never demonstrated that microscopic endometriosis, whether in the peritoneum or in lymph nodes (and maybe in the bowel), and also subtle lesions are a cause of pain or infertility. It is at least surprising that progression of these lesions still is postulated without being demonstrated except *in vitro* [56]. The traditional concepts of pathophysiology cannot explain all the different types of endometriosis [55] and do not explain heredity

and the fact that cystic and deep endometriosis are clonal in origin. A genomic mutation of these cells [135] could explain why and when glands and stroma outside the uterus are normal or pathologic. It would facilitate understanding clonality and heredity. It could explain why lesions develop into typical, cystic, and deep lesions. It could explain that typical, cystic, and deep lesions are end stages—similar to a no longer growing myoma—without progression from one type to another as clinical observation suggests. It also could explain why in baboons and the rhesus monkeys induction of severe lesions has failed. It is consistent with primate data of the heredity of spontaneous typical lesions [136], with the fact that total body irradiation [3] induces typical endometriotic lesions and that dioxin induces cystic ovarian endometriosis, the severity of which was dose dependent [1]. These data are all consistent with the fact that it takes many years before a genomic incident becomes apparent [1, 137]. Recently, lesions similar to deep endometriosis were found by injecting slices of endometrium with myometrium [116] pointing to a relationship between functional endometrium, basal endometrium, and junctional zone. The role of the junctional zone only recently begins to receive attention.

A fundamental question is whether cells in subtle, typical, cystic, and deep lesions are genetically identical or different from endometrial cells. Similarly, it is important to understand whether the differences of endometrium in women with and without endometriosis are the consequence of endometriosis or are a sign of vulnerability and of predisposition to develop disease. In the absence of data, it seems prudent today to replace endometriosis by subtle endometriosis, typical endometriosis, cystic endometriosis, and deep endometriosis. This would highlight that they are potentially different. Typical, cystic, and deep endometriosis could be considered as a benign tumor similar to a myoma or a polyp generally having a genomic alteration, facilitated by the genetic background, and epigenetic modifications [138]. It would facilitate to understand that some deep endometriosis lesions are different and clinically fast progressive as suggested by clinical observation (personal observations) and evidenced by complications during pregnancy [89].

Considering subtle, typical, cystic, and deep endometriosis as separate entities changes our view on epidemiology. There is no hard evidence for an increase in the prevalence of typical, cystic, or deep endometriosis or for a specific causal factor. What remains is that the prevalences of typical, cystic, and deep endometriosis increase with age which is consistent with the observation that the prevalence of most benign tumors increases with age. Considering these lesions as benign tumors on the other hand risks to increase our concern of a relationship with dioxin, PCB [2], and

radiation, which all act at the level of the genome. Genetic predisposition moreover is easily connected with cancer development. Although weak as evidence, the impression by authors and other surgeons who all are pioneers of deep endometriosis surgery with a large personal experience that deep lesions become more severe and probably more frequent over the last 20 years should not be disregarded.

Considering subtle, typical, cystic, and deep endometriosis as separate entities could change the design of epidemiologic studies. Since deep endometriosis is associated with very severe pain in most women, the majority will undergo surgery. If considered as adenomyosis externa, diagnosis by pathology will carry little errors and the volume of the lesion, a measure of severity, will be available. Since recurrence rates of deep endometriosis are small, even women that undergo repetitive surgery can be identified and anyway should cause little bias. Even in women that undergo hysterectomy, the diagnosis of deep endometriosis can be made. This should permit hospital discharge record-based case-control studies, with as only bias that not all women undergo surgery, some 5 % being pain free. Epidemiology of deep endometriosis should permit to evaluate an eventual relationship with pollution, radioactivity, and our changing lifestyle with a delay in childbirth. Epidemiology of deep endometriosis will in addition permit to evaluate questions such as whether the use of oral contraception and the years of medical treatment for pelvic pain without a diagnosis might stimulate the development of some severe forms of endometriosis.

We should learn from the past, especially from the excellent available data of epidemiology [4–6] and associated diseases such as acne [7–12] or with food intake as fat consumption [100, 101] of endometriosis. Studies of the epidemiology of subtle, typical, cystic, and deep endometriosis would eliminate the question whether microscopical and subtle lesions should be considered pathology. The design would fully comply with the conclusion [13] that to be pathologic, endometrial glands and stroma should be progressive and invasive and that this “may increase the likelihood of observing true associations in etiologic studies” and avoid “threats to validity of substantial magnitude that exist in both clinic-based and population-based epidemiologic studies of endometriosis.”

In conclusion, subtle, typical, cystic, and deep endometriosis lesions should be considered as separate entities, until pathophysiology and natural history are known. This would permit to answer whether subtle lesions are pathologic and whether and which lesions are recurrent and progressive. It could answer the question whether endometrial changes in endometriosis are the hallmark of pre-existing susceptibility. This would permit to run epidemiologic studies on deep endometriosis with little bias, which will permit to answer the question whether prevalence is increasing or not. This ultimately would permit to answer the questions whether pollution, radiation, and lifestyle would be etiologically involved.

**Acknowledgments** We thank Jacques Donnez, Brussels, Belgium; Antonio Setubal, Lisbon Portugal; Charles Koh, Milwaukee, USA; and Camran, Ceana, and Farr Nezhat, Atlanta, USA, for their clinical opinion on prevalence and severity of deep endometriosis over the last 20 years.

**Compliance with ethical standards**

**Funding** The authors did not have any funding.

**Conflict of interest** Neither Koninckx, nor Ussia, nor Keckstein, nor Wattiez, nor Adamyan report a conflict of interest for this study. Koninckx is shareholder of Endosat.

**Ethical approval** Not applicable.

**Informed consent** Not applicable.

**Statement of authorship** All authors are recognized deep endometriosis surgeons, active over the last 20 years with a large personal experience. The manuscript was initiated and conceived in many preparatory discussions, written by PK, and all other authors contributed in reviewing the data and the manuscript.

## References

1. Rier SE, Martin DC, Bowman RE, Dmowski WP, Becker JL (1993) Endometriosis in rhesus monkeys (*Macaca mulatta*) following chronic exposure to 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Fundam Appl Toxicol* 21:433–441
2. Bimbaum LS, Cummings AM (2002) Dioxins and endometriosis: a plausible hypothesis. *Environ Health Perspect* 110:15–21
3. Fanton JW, Golden JG (1991) Radiation-induced endometriosis in *Macaca mulatta*. *Radiat Res* 126:141–146
4. Eskenazi B, Warner ML (1997) Epidemiology of endometriosis. *Obstet Gynecol Clin N Am* 24:235–258
5. Vigano P, Parazzini F, Somigliana E, Vercellini P (2004) Endometriosis: epidemiology and aetiological factors. *Best Pract Res Clin Obstet Gynaecol* 18:177–200
6. Missmer SA, Cramer DW (2003) The epidemiology of endometriosis. *Obstet Gynecol Clin N Am* 30:1–19 vii
7. Harris HR, Costenbader KH, Mu F, Kvaskoff M, Malspeis S, Karlson EW et al (2015) Endometriosis and the risks of systemic lupus erythematosus and rheumatoid arthritis in the Nurses' Health Study II. *Ann Rheum Dis*
8. Kvaskoff M, Mu F, Terry KL, Harris HR, Poole EM, Farland L, et al. (2015) Endometriosis: a high-risk population for major chronic diseases? *Hum Reprod Update* 21:500–516
9. Missmer SA (2015) Safety in reproductive medicine: breadth, depth and discovery. *Hum Reprod* 30:2252–2253
10. Mahalingaiah S, Hart JE, Laden F, Aschengrau A, Missmer SA (2014) Air pollution exposures during adulthood and risk of endometriosis in the Nurses' health study II. *Environ Health Perspect* 122:58–64
11. Xie J, Kvaskoff M, Li Y, Zhang M, Qureshi AA, Missmer SA, et al. (2014) Severe teenage acne and risk of endometriosis. *Hum Reprod* 29:2592–2599
12. Shah DK, Missmer SA (2011) Scientific investigation of endometriosis among adolescents. *J Pediatr Adolesc Gynecol* 24:S18–S19
13. Holt VL, Weiss NS (2000) Recommendations for the design of epidemiologic studies of endometriosis. *Epidemiology* 11: 654–659



14. Cullen TS (1896) Adeno-myoma of the round ligament. *Johns Hopkins Hosp Bull* 7:112–114
15. Cullen TS (1896) Adenoma-myoma uteri diffusum benignum. *J Hopkins Hosp Bull* 6:133–137
16. Cullen TS (1919) The distribution of adenomyomata containing uterine mucosa. *Am J Obstet Gynecol* 80:130–138
17. Meyer R (1919) Über den stand der frageder Adenomyositis und Adenomyome im Allgemeine und ins Besondere über Adenomyositis seroepithelialis und Adenomyometritis sarcomatosa. *Zentralbl Gynakol* 36:745
18. Meyer R (1923) Zur frage der Urnieren-genese van Adenomyomen. *Zentralbl Gynakol* 15:577–587
19. Sampson JA (1925) Heterotopic or misplaced endometrial tissue. *Am J Obstet Gynecol* 10:649–664
20. Sampson JA (1927) Peritoneal endometriosis due to the menstrual dissemination of endometrial tissue into the peritoneal cavity. *Am J Obstet Gynecol* 14:422–469
21. Scott RB, Telinde RW (1950) External endometriosis—the scourge of the private patient. *Ann Surg* 131:697
22. Burgmeister RE, Fechner RE, Franklin RR (1969) Endosalpingiosis of the peritoneum. *Obstet Gynecol* 34:310–317
23. Latcher JW (1953) Endometriosis of the umbilicus. *Am J Obstet Gynecol* 66:161
24. Brosens IA, Koninckx PR, Corveleyn PA (1978) A study of plasma progesterone, oestradiol-17beta, prolactin and LH levels, and of the luteal phase appearance of the ovaries in patients with endometriosis and infertility. *Br J Obstet Gynaecol* 85:246–250
25. Jansen RP (1986) Minimal endometriosis and reduced fecundability: prospective evidence from an artificial insemination by donor program. *Fertil Steril* 46:141–143
26. Stripling MC, Martin DC, Chatman DL, Vander Zwaag R, Poston WM (1988) Subtle appearance of pelvic endometriosis. *Fertil Steril* 49:427–431
27. Martin DC, Ahmic R, El Zeky FA, Vander Zwaag R, Pickens MT, Cherry K (1990) Increased histologic confirmation of endometriosis. *J Gynecol Surg* 6:275–279
28. Martin DC, Hubert GD, Vander ZR, El Zeky FA (1989) Laparoscopic appearances of peritoneal endometriosis. *Fertil Steril* 51:63–67
29. Walter AJ, Hentz JG, Magtibay PM, Cornella JL, Magrina JF (2001) Endometriosis: correlation between histologic and visual findings at laparoscopy. *Am J Obstet Gynecol* 184:1407–1411
30. Moen MH, Halvorsen TB (1992) Histologic confirmation of endometriosis in different peritoneal lesions. *Acta Obstet Gynecol Scand* 71:337–342
31. Von Rokitanski C (1860) Über Uterusdrüsen-neubildung in uterus und Ovarial-Sarcomen. *Zeitschrift GMS Ärzte Wien* 37:577–581
32. Russell WW (1899) Aberrant portions of the mullerian duct found in an ovary. *Johns Hopkins Hosp Bull* 94-96:8–10
33. Jimbo H, Hitomi Y, Yoshikawa H, Yano T, Momoeda M, Sakamoto A, et al. (1997) Evidence for monoclonal expansion of epithelial cells in ovarian endometrial cysts. *Am J Pathol* 150:1173–1178
34. Sampson JA (1940) The development of the implantation theory for the development of endometriosis. *Am J Obstet Gynecol* 40:549–557
35. Gruenwald P (1942) Origin of endometriosis from the mesenchyme of the celomic walls. *Am J Obstet Gynecol* 44:470–474
36. Koninckx PR, Ide P, Vandenbroucke W, Brosens IA (1980) New aspects of the pathophysiology of endometriosis and associated infertility. *J Reprod Med* 24:257–260
37. Halme J, Hammond MG, Hulka JF, et al. (1984) Retrograde menstruation in healthy women and in patients with endometriosis. *Obstet Gynecol* 64:151–154
38. Kruiwagen RFFM, Poels LG, Willemsen WNP, De Ronde IJY, Jap PHK, Rolland R (1991) Endometrial epithelial cells in peritoneal fluid during the early follicular phase. *Fertil Steril* 55:297–303
39. Vasquez G, Comillie F, Brosens IA (1984) Peritoneal endometriosis: scanning electron microscopy and histology of minimal pelvic endometriotic lesions. *Fertil Steril* 42:696–703
40. Murphy AA, Green WR, Bobbie D, dela Cruz ZC, Rock JA (1986) Unsuspected endometriosis documented by scanning electron microscopy in visually normal peritoneum. *Fertil Steril* 46:522–524
41. Murphy AA, Guzick DS, Rock JA (1989) Microscopic peritoneal endometriosis [letter; comment]. *Fertil Steril* 51:1072–1074
42. Nisolle M, Paindaveine B, Bourdon A, Berliere M, Casanas Roux F, Donnez J (1990) Histologic study of peritoneal endometriosis in infertile women. *Fertil Steril* 53:984–988
43. Balasch J, Creus M, Fabregues F, Carmona F, Ordi J, Martinez-Roman S, et al. (1996) Visible and non-visible endometriosis at laparoscopy in fertile and infertile women and in patients with chronic pelvic pain: a prospective study. *Hum Reprod* 11:387–391
44. Khan KN, Fujishita A, Kitajima M, Hiraki K, Nakashima M, Masuzaki H (2014) Occult microscopic endometriosis: undetectable by laparoscopy in normal peritoneum. *Hum Reprod* 29:462–472
45. Koninckx PR, Meuleman C, Demeyere S, Lesaffre E, Cornillie FJ (1991) Suggestive evidence that pelvic endometriosis is a progressive disease, whereas deeply infiltrating endometriosis is associated with pelvic pain. *Fertil Steril* 55:759–765
46. Koninckx PR (1994) Is mild endometriosis a condition occurring intermittently in all women? *Hum Reprod* 9:2202–2205
47. Evers JL, Dunselman GA, Groothuis P (2005) Now you see them, now you don't. *Fertil Steril* 84:31–32
48. Rossini R, Monsellato D, Bertolaccini L, Pesci A, Zamboni G, Ceccaroni M et al (2016) Lymph nodes involvement in deep infiltrating intestinal endometriosis: does it really mean anything? *J Minim Invasive Gynecol*
49. Koninckx PR (1998) Biases in the endometriosis literature. Illustrated by 20 years of endometriosis research in Leuven. *Eur J Obstet Gynecol Reprod Biol* 81:259–271
50. Mechsner S, Weichbrodt M, Riedlinger WF, Kaufmann AM, Schneider A, Kohler C (2010) Immunohistochemical evaluation of endometriotic lesions and disseminated endometriosis-like cells in incidental lymph nodes of patients with endometriosis. *Fertil Steril* 94:457–463
51. Badescu A, Roman H, Aziz M, Puscasiu L, Molnar C, Huet E, et al. (2016) Mapping of bowel occult microscopic endometriosis implants surrounding deep endometriosis nodules infiltrating the bowel. *Fertil Steril* 105:430–434
52. Roman H, Henrietier C, Darwish B, Badescu A, Csanyi M, Aziz M, et al. (2016) Bowel occult microscopic endometriosis in resection margins in deep colorectal endometriosis specimens has no impact on short-term postoperative outcomes. *Fertil Steril* 105:423–429
53. Koninckx PR, Donnez J, Brosens I (2015) Microscopic endometriosis: impact on our understanding of the disease and its surgery. *Fertil Steril*
54. Koninckx PR, Oosterlynck D, D'Hooghe T, Meuleman C (1994) Deeply infiltrating endometriosis is a disease whereas mild endometriosis could be considered a non-disease. *Ann N Y Acad Sci* 734:333–341
55. Vercellini P, Vigano P, Somigliana E, Fedele L (2014) Endometriosis: pathogenesis and treatment. *Nat Rev Endocrinol* 10:261–275
56. Nap AW, Groothuis PG, Demir AY, Evers JL, Dunselman GA (2004) Pathogenesis of endometriosis. *Best Pract Res Clin Obstet Gynaecol* 18:233–244
57. Olive DL, Henderson DY (1987) Endometriosis and mullerian anomalies. *Obstet Gynecol* 69:412–415

58. Adamyan LV, Kulakov VI, Hashukoeva AZ (1998) Malformations of the uterus and vagina. In: Adamyan LV, Kulakov VI (eds) Endometriosis in medicine edited by Moscow medicina, Oformlenie Antidor
59. Adamyan LV, Yarotskaya EL (2007) Endoscopic diagnosis and correction of malformations of female genitalia. In: Pasic RP, Levine RL (eds) A Practical Manual of Laparoscopy and Minimally Invasive Gynecology: A Clinical Cookbook. second edition ed. Informa Healthcare, New York. 347–364
60. Tong J, Zhu L, Chen N, Lang J (2014) Endometriosis in association with Herlyn-Werner-Wunderlich syndrome. *Fertil Steril* 102:790–794
61. Koninckx PR, Kennedy SH, Barlow DH (1998) Endometriotic disease: the role of peritoneal fluid. *Hum Reprod Update* 4:741–751
62. Kennedy SH, Mardon H, Barlow DH (1995) Familial endometriosis. *J Assist Reprod Genet* 12:32–34
63. Kennedy S, Hadfield R, Mardon H, Barlow D (1996) Age of onset of pain symptoms in non-twin sisters concordant for endometriosis. *Hum Reprod* 11:403–405
64. Hadfield RM, Mardon HJ, Barlow DH, Kennedy SH (1997) Endometriosis in monozygotic twins. *Fertil Steril* 68:941–942
65. Hadfield RM, Manek S, Nakago S, Mukherjee S, Weeks DE, Mardon HJ, et al. (1999) Absence of a relationship between endometriosis and the N314D polymorphism of galactose-1-phosphate uridyl transferase in a UK population. *Mol Hum Reprod* 5:990–993
66. Hadfield RM, Manek S, Weeks DE, Mardon HJ, Barlow DH, Kennedy SH (2001) Linkage and association studies of the relationship between endometriosis and genes encoding the detoxification enzymes GSTM1, GSTT1 and CYP1A1. *Mol Hum Reprod* 7:1073–1078
67. Hadfield RM, Pullen JG, Davies KF, Wolfensohn SE, Kemnitz JW, Weeks DE, et al. (2001) Toward developing a genome-wide microsatellite marker set for linkage analysis in the rhesus macaque (*Macaca mulatta*): identification of 76 polymorphic markers. *Am J Primatol* 54:223–231
68. Nakago S, Hadfield RM, Zondervan KT, Mardon H, Manek S, Weeks DE, et al. (2001) Association between endometriosis and N-acetyl transferase 2 polymorphisms in a UK population. *Mol Hum Reprod* 7:1079–1083
69. Kennedy S (2003) Genetics of endometriosis: a review of the positional cloning approaches. *Semin Reprod Med* 21:111–118
70. Yano T, Jimbo H, Yoshikawa H, Tsutsumi O, Taketani Y (1999) Molecular analysis of clonality in ovarian endometrial cysts. *Gynecol Obstet Investig* 47(Suppl 1):41–45
71. Tamura M, Fukaya T, Murakami I, Uehara S, Yajima A (1998) Analysis of clonality in human endometriotic cysts based on evaluation of X chromosome inactivation in archival formalin-fixed, paraffin-embedded tissue. *Lab Invest* 78:213–218
72. Koninckx PR (1999) The physiopathology of endometriosis: pollution and dioxin. *Gynecol Obstet Investig* 47(Suppl 1):47–49
73. Dmowski WP (1992) Immunology of endometriosis. *Arq Matern Dr Alfredo Da Costa* 40–43
74. Berkkanoglu M, Arici A (2003) Immunology and endometriosis. *Am J Reprod Immunol* 50:48–59
75. Koninckx PR, Kennedy SH, Barlow DH (1999) Pathogenesis of endometriosis: the role of peritoneal fluid. *Gynecol Obstet Investig* 47(Suppl 1):23–33
76. Oosterlynck DJ, Cornillie FJ, Waer M, Vandeputte M, Koninckx PR (1991) Women with endometriosis show a defect in natural killer activity resulting in a decreased cytotoxicity to autologous endometrium. *Fertil Steril* 56:45–51
77. Oosterlynck DJ, Meuleman C, Waer M, Koninckx PR, Vandeputte M (1993) Immunosuppressive activity of peritoneal fluid in women with endometriosis. *Obstet Gynecol* 82:206–212
78. Oosterlynck DJ, Meuleman C, Waer M, Koninckx PR (1994) CO<sub>2</sub>-laser excision of endometriosis does not improve the decreased natural killer activity. *Acta Obstet Gynecol Scand* 73: 333–337
79. Ibrahim MG, Chiantera V, Frangini S, Younes S, Kohler C, Taube ET, et al. (2015) Ultramicro-trauma in the endometrial-myometrial junctional zone and pale cell migration in adenomyosis. *Fertil Steril* 104:1475–1483
80. Brosens I, Robertson WB, Dixon HG (1967) The physiological response of the vessels of the placental bed to normal pregnancy. *J Pathol Bacteriol* 93:569–579
81. Brosens IA, Robertson WB, Dixon HG (1972) The role of the spiral arteries in the pathogenesis of preeclampsia. *Obstet Gynecol Annu* 1:177–191
82. Brosens I, Curcic A, Vejnovic T, Gargett CE, Brosens JJ, Benagiano G (2015) The perinatal origins of major reproductive disorders in the adolescent: research avenues. *Placenta* 36: 341–344
83. Vigano P, Corti L, Berlanda N (2015) Beyond infertility: obstetrical and postpartum complications associated with endometriosis and adenomyosis. *Fertil Steril*
84. Aris A (2014) A 12-year cohort study on adverse pregnancy outcomes in eastern townships of Canada: impact of endometriosis. *Gynecol Endocrinol* 30:34–37
85. Brosens JJ, Parker MG, McIndoe A, Pijnenborg R, Brosens IA (2009) A role for menstruation in preconditioning the uterus for successful pregnancy. *Am J Obstet Gynecol*
86. Nap AW, Dunselman GA, Griffioen AW, Mayo KH, Evers JL, Groothuis PG (2005) Angiostatic agents prevent the development of endometriosis-like lesions in the chicken chorioallantoic membrane. *Fertil Steril* 83:793–795
87. Busacca M, Marana R, Caruana P, Candiani M, Muzii L, Calia C, et al. (1999) Recurrence of ovarian endometrioma after laparoscopic excision. *Am J Obstet Gynecol* 180:519–523
88. Muzii L, Miller CE (2011) The singer, not the song. *J Minim Invasive Gynecol* 18:666–667
89. Setubal A, Sidiropoulou Z, Torgal M, Casal E, Lourenco C, Koninckx P (2014) Bowel complications of deep endometriosis during pregnancy or in vitro fertilization. *Fertil Steril* 101: 442–446
90. Koninckx PR, Muyltermans M, Moerman P, Meuleman C, Deprest J, Cornillie F (1992) CA 125 concentrations in ovarian 'chocolate' cyst fluid can differentiate an endometriotic cyst from a cystic corpus luteum. *Hum Reprod* 7:1314–1317
91. Brosens IA (1994) Endoscopic exploration and classification of the chocolate cysts. *Hum Reprod* 9:2213–2214
92. Martin DC, Demos BJ (1990) Histology of chocolate cysts. *J Gynecol Surg* 6:43–46
93. Brosens IA, Puttemans PJ, Deprest J (1994) The endoscopic localization of endometrial implants in the ovarian chocolate cyst. *Fertil Steril* 61:1034–1038
94. Mais V, Guerriero S, Ajossa S, Angiolucci M, Paoletti AM, Melis GB (1993) The efficiency of transvaginal ultrasonography in the diagnosis of endometrioma. *Fertil Steril* 60:776–780
95. Van HC, Van CB, Guerriero S, Savelli L, Paladini D, Lissoni AA, et al. (2010) Endometriomas: their ultrasound characteristics. *Ultrasound Obstet Gynecol* 35:730–740
96. Gordts S, Puttemans P, Gordts S, Valkenburg M, Brosens I, Campo R (2014) Transvaginal endoscopy and small ovarian endometriomas: unravelling the missing link? *Gynecol Surg* 11: 3–7
97. Koninckx PR, Meuleman C, Oosterlynck D, Cornillie FJ (1996) Diagnosis of deep endometriosis by clinical examination during menstruation and plasma CA-125 concentration. *Fertil Steril* 65: 280–287
98. Cornillie FJ, Oosterlynck D, Lauweryns JM, Koninckx PR (1990) Deeply infiltrating pelvic endometriosis: histology and clinical significance. *Fertil Steril* 53:978–983

99. Koninckx PR, Martin DC (1992) Deep endometriosis: a consequence of infiltration or retraction or possibly adenomyosis externa? *Fertil Steril* 58:924–928
100. Koninckx PR, Brosens IA (2011) Dietary fat consumption and endometriosis risk. *Hum Reprod* 26:731–732
101. Missmer SA, Chavarro JE, Malspeis S, Bertone-Johnson ER, Hornstein MD, Spiegelman D, et al. (2010) A prospective study of dietary fat consumption and endometriosis risk. *Hum Reprod* 25:1528–1535
102. Spitsyn VA, Andreeva EN, Adamyan LV, Agapova RK, Sandoval KR (1996) Genetic aspects of endometriosis: peculiarities of the distribution of polymorphic gene frequency. *Genetika* 32:1693–1699
103. Rahmioglu N, Nyholt DR, Morris AP, Missmer SA, Montgomery GW, Zondervan KT (2014) Genetic variants underlying risk of endometriosis: insights from meta-analysis of eight genome-wide association and replication datasets. *Hum Reprod Update* 20:702–716
104. Foster WG (2008) Endocrine toxicants including 2,3,7,8-tetrachlorodibenzo-p-dioxin (TCDD) and dioxin-like chemicals and endometriosis: is there a link? *J Toxicol Environ Health B Crit Rev* 11:177–187
105. Montgomery GW, Nyholt DR, Zhao ZZ, Treloar SA, Painter JN, Missmer SA, et al. (2008) The search for genes contributing to endometriosis risk. *Hum Reprod Update* 14:447–457
106. Heilier JF, Donnez J, Lison D (2008) Organochlorines and endometriosis: a mini-review. *Chemosphere* 71:203–210
107. Jimbo H, Yoshikawa H, Onda T, Yasugi T, Sakamoto A, Taketani Y (1997) Prevalence of ovarian endometriosis in epithelial ovarian cancer. *Int J Gynaecol Obstet* 59:245–250
108. Redwine DB (1987) Age-related evolution in color appearance of endometriosis. *Fertil Steril* 48:1062–1063
109. Vercellini P, Parazzini F, Bolis G, Carinelli S, Dindelli M, Vendola N, et al. (1993) Endometriosis and ovarian cancer. *Am J Obstet Gynecol* 169:181–182
110. Laufer MR, Sanfilippo J, Rose G (2003) Adolescent endometriosis: diagnosis and treatment approaches. *J Pediatr Adolesc Gynecol* 16:S3–11
111. Cramer DW, Missmer SA (2002) The epidemiology of endometriosis. *Ann N Y Acad Sci* 955:11–22
112. Moen MH, Schei B (1997) Epidemiology of endometriosis in a Norwegian county. *Acta Obstet Gynecol Scand* 76:559–562
113. Matorras R, Rodriguez F, Pijoan JI, Etxanojauregui A, Neyro JL, Elorriaga MA, et al. (2001) Women who are not exposed to spermatozoa and infertile women have similar rates of stage I endometriosis. *Fertil Steril* 76:923–928
114. Janssen EB, Rijkers AC, Hoppenbrouwers K, Meuleman C, D'Hooghe TM (2013) Prevalence of endometriosis diagnosed by laparoscopy in adolescents with dysmenorrhea or chronic pelvic pain: a systematic review. *Hum Reprod Update* 19:570–582
115. Guo SW, Wang Y (2006) The prevalence of endometriosis in women with chronic pelvic pain. *Gynecol Obstet Investig* 62:121–130
116. Donnez O, Van LA, Defrere S, Colette S, Van KO, Dehoux JP et al (2012) Induction of endometriotic nodules in an experimental baboon model mimicking human deep nodular lesions. *Fertil Steril*
117. Rier SE, Turner WE, Martin DC, Morris R, Lucier GW, Clark GC (2001) Serum levels of TCDD and dioxin-like chemicals in rhesus monkeys chronically exposed to dioxin: correlation of increased serum PCB levels with endometriosis. *Toxicol Sci* 59:147–159
118. Koninckx PR, Braet P, Kennedy SH, Barlow DH (1994) Dioxin pollution and endometriosis in Belgium. *Hum Reprod* 9:1001–1002
119. Eskenazi B, Mocarelli P, Warner M, Samuels S, Vercellini P, Olive D, et al. (2002) Serum dioxin concentrations and endometriosis: a cohort study in Seveso, Italy. *Environ Health Perspect* 110:629–634
120. Tsutsumi O, Momoeda M, Takai Y, Ono M, Taketani Y (2000) Breast-fed infants, possibly exposed to dioxins in milk, have unexpectedly lower incidence of endometriosis in adult life. *Int J Gynaecol Obstet* 68:151–153
121. Braun DP, Ding J, Dmowski WP (2002) Peritoneal fluid-mediated enhancement of eutopic and ectopic endometrial cell proliferation is dependent on tumor necrosis factor-alpha in women with endometriosis. *Fertil Steril* 78:727–732
122. Ding J, Shen J, Braun DP, Rana N, Dmowski WP (2002) Endometrial apoptosis, macrophage content, and TNFalpha expression in endometriosis. *Fertil Steril* 77(Suppl 1):S1–S2
123. Oosterlynck, D (1983) PhD thesis: natural killer cells and endometriosis
124. Oosterlynck DJ, Cornillie FJ, Waer M, Koninckx PR (1993) Immunohistochemical characterization of leucocyte subpopulations in endometriotic lesions. *Arch Gynecol Obstet* 253:197–206
125. Oosterlynck DJ, Meuleman C, Waer M, Koninckx PR (1994) Transforming growth factor-beta activity is increased in peritoneal fluid from women with endometriosis. *Obstet Gynecol* 83:287–292
126. Oosterlynck DJ, Lacquet FA, Waer M, Koninckx PR (1994) Lymphokine-activated killer activity in women with endometriosis. *Gynecol Obstet Investig* 37:185–190
127. Oosterlynck DJ, Meuleman C, Waer M, Vandeputte M, Koninckx PR (1992) The natural killer activity of peritoneal fluid lymphocytes is decreased in women with endometriosis. *Fertil Steril* 58:290–295
128. Vinatier D, Dufour P, Oosterlynck D (1996) Immunological aspects of endometriosis. *Hum Reprod Update* 2:371–384
129. Koninckx PR (1987) Pelvic endometriosis: a consequence of stress? *Contrib Gynecol Obstet* 16:56–59
130. Demyttenaere K, Nijs P, Evers-Kiebooms G, Koninckx PR (1991) Coping, ineffectiveness of coping and the psychoendocrinological stress responses during in-vitro fertilization. *J Psychosom Res* 35:231–243
131. Demyttenaere K, Nijs P, Evers-Kiebooms G, Koninckx PR (1989) The effect of a specific emotional stressor on prolactin, cortisol, and testosterone concentrations in women varies with their trait anxiety. *Fertil Steril* 52:942–948
132. Parazzini F, Luchini L, Vezzoli F, Mezzanotte C, Vercellini P, Romanini C, et al. (1994) Prevalence and anatomical distribution of endometriosis in women with selected gynaecological conditions: results from a multicentric Italian study. *Hum Reprod* 9:1158–1162
133. Mahmood TA, Templeton A (1991) Prevalence and genesis of endometriosis. *Hum Reprod* 6:544–549
134. Koninckx PR, Batt RE, Hummelshoj L, McVeigh E, Ussia A, Yeh J (2010) The elephant in the room: quality control of endometriosis data. *J Minim Invasive Gynecol* 17:637–640
135. Koninckx PR, Barlow D, Kennedy S (1999) Implantation versus infiltration: the Sampson versus the endometriotic disease theory. *Gynecol Obstet Investig* 47(Suppl 1):3–9
136. Zondervan K, Cardon L, Desrosiers R, Hyde D, Kemnitz J, Mansfield K, et al. (2002) The genetic epidemiology of spontaneous endometriosis in the rhesus monkey. *Ann N Y Acad Sci* 955:233–238
137. Rier SE (2008) Environmental immune disruption: a comorbidity factor for reproduction? *Fertil Steril* 89:e103–e108
138. Kobayashi H, Imanaka S, Nakamura H, Tsuji A (2014) Understanding the role of epigenomic, genomic and genetic alterations in the development of endometriosis (review). *Mol Med Rep* 9:1483–1505